


Gravitational Waves


Supplement to Lecture XV

October 23, 2019

LIGO – Hanford Site (1 of 2)



Advanced LIGO Optical Layout of the Interferometer





Observation of Gravitational Waves from a Binary Black Hole Merger

B. P. Abbott *et al.**

(LIGO Scientific Collaboration and Virgo Collaboration)

(Received 21 January 2016; published 11 February 2016)

On September 14, 2015 at 09:50:45 UTC the two detectors of the Laser Interferometer Gravitational-Wave Observatory simultaneously observed a transient gravitational-wave signal. The signal sweeps upwards in frequency from 35 to 250 Hz with a peak gravitational-wave strain of 1.0×10^{-21} . It matches the waveform predicted by general relativity for the inspiral and merger of a pair of black holes and the ringdown of the resulting single black hole. The signal was observed with a matched-filter signal-to-noise ratio of 24 and a false alarm rate estimated to be less than 1 event per 203 000 years, equivalent to a significance greater than 5.1σ . The source lies at a luminosity distance of 410_{-180}^{+160} Mpc corresponding to a redshift $z = 0.09_{-0.04}^{+0.03}$. In the source frame, the initial black hole masses are $36_{-4}^{+5} M_{\odot}$ and $29_{-4}^{+4} M_{\odot}$, and the final black hole mass is $62_{-4}^{+4} M_{\odot}$, with $3.0_{-0.5}^{+0.5} M_{\odot} c^2$ radiated in gravitational waves. All uncertainties define 90% credible intervals. These observations demonstrate the existence of binary stellar-mass black hole systems. This is the first direct detection of gravitational waves and the first observation of a binary black hole merger.

DOI: [10.1103/PhysRevLett.116.061102](https://doi.org/10.1103/PhysRevLett.116.061102)





TABLE I. Source parameters for GW150914. We report median values with 90% credible intervals that include statistical errors, and systematic errors from averaging the results of different waveform models. Masses are given in the source frame; to convert to the detector frame multiply by $(1+z)$ [90]. The source redshift assumes standard cosmology [91].



Primary black hole mass	$36_{-4}^{+5} M_{\odot}$
Secondary black hole mass	$29_{-4}^{+4} M_{\odot}$
Final black hole mass	$62_{-4}^{+4} M_{\odot}$
Final black hole spin	$0.67_{-0.07}^{+0.05}$
Luminosity distance	410_{-180}^{+160} Mpc
Source redshift z	$0.09_{-0.04}^{+0.03}$

The estimated total energy radiated in gravitational waves is $3.0_{-0.5}^{+0.5} M_{\odot} c^2$. The system reached a peak gravitational-wave luminosity of $3.6_{-0.4}^{+0.5} \times 10^{56}$ erg/s, equivalent to $200_{-20}^{+30} M_{\odot} c^2/s$.




GW 170817

(LIGO + Virgo Collaborations 2017)



$$m_T^{2/5} \mu^{3/5} = 1.188_{-0.002}^{+0.004} M_{\text{Sun}}$$




Gamma Ray Burst Coincident with GW Event

(LIGO, Virgo, Fermi-GBM,
INTEGRAL Collaborations,
2017)

Multi-Wavelength Electromagnetic Observations of GW170817

(Many Collaborations, 2017)



Host galaxy is NGC 4993, $z = 0.0097$

H_0 from 1 Gravitational Wave Source

